

1D NMR Spectra do not work in QSPR



Published January 27, 2006

Citation

Willighagen, E. (2006, January 27). 1D NMR Spectra do not work in QSPR. *Chem-bla-ics*. <https://doi.org/10.59350/h25t3-mpk14>

Keywords

Cheminf

Copyright

Copyright © None 2006. Distributed under the terms of the [Creative Commons Attribution 4.0 International License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

About two years ago a student started with me to work on the use of 1D NMR and IR spectra in quantitative structure-activity relationship (QSAR) work, with the goal to show that these spectra contain 3D information relevant to QSAR models. It is known that these spectra depend on the 3D conformation of the molecule.

Half a year later we concluded that from the data which we started with (48 compounds with binding affinity), no conclusions could be drawn what so ever: no statistically sound models could be build at all. So, we composed three larger data sets. These sets, all QSPR data sets, did give us models, but all the spectra based models were worse than a [Dragon](#) descriptor based model using the same number of variables, without doing any variable selection.

I presented this work at the 7th [ICCS](#) in Noordwijkerhout half a year ago, and now got published in the JCI: DOI [10.1021/ci050282s](https://doi.org/10.1021/ci050282s). Comments on this article are **most** welcome!